

References CBDA How Can it Help?

- (1) Pellesi, Lanfranco, et al. " European Journal of Clinical Pharmacology 74.11 (2018): 1427-1436.
- (2) S Takeda et al. Cannabidiolic Acid as a Selective Cyclooxygenase-2 Inhibitory Component in Cannabis. Drug Metabolism and Disposition September 2008, 36 (9) 1917-1921
- (3) EM Rock et al. Effect of low doses of cannabidiolic acid and ondansetron on LiCl-induced conditioned gaping (a model of nausea-induced behaviour) in rats. British journal of pharmacology vol. 169,3 (2013): 685-92.
- (4) Rock, Erin M., et al. Psychopharmacology 234.14 (2017): 2207-2217
- (5) Anderson, Lyndsey L., et al. Journal of natural products 82.11 (2019): 3047-3055.
- (6) R Mechoulam et al. Hashish. IV. The isolation and structure of cannabinolic cannabidiolic and cannabigerolic acids. Tetrahedron 1965 May;21(5):1223-9
- (7) EM Rock et al. Cannabidiolic acid prevents vomiting in Suncus murinus and nausea-induced behaviour in rats by enhancing 5-HT1A receptor activation. Br J Pharmacol. 2013 Mar; 168(6): 1456–1470
- (8) Use of cannabinoids in the treatment of epilepsy. Patent application. GW Pharma
- (9) S Takeda et al. Down-regulation of cyclooxygenase-2 (COX-2) by cannabidiolic acid in human breast cancer cells. The Journal of Toxicological Sciences. Vol.39, No.5, 711-716, 2014

(10) D Hen-Shoval et al. Acute oral cannabidiolic acid methyl ester reduces depression-like behavior in two genetic animal models of depression. Behav Brain Res 2018 Oct 1;351:1-3